

ATOMIC PHYSICS: COMPUTER CALCULATIONS AND THEORETICAL ANALYSIS

E. G. Drukarev

*Petersburg Nuclear Physics Institute,
St. Petersburg, Gatchina 188300, Russia*

Abstract

It is demonstrated, how the theoretical analysis preceding the numerical calculations helps to calculate the energy of the ground state of helium atom, and enables to avoid qualitative errors in calculations of the characteristics of double photoionization.

1 Introduction

As it stands now, many of the publications devoted to interaction of photons and electrons with atoms and molecules have similar structure. In the first step a most general quantum mechanics formula for the cross section is written. In the next step the most "accurate" numerical functions for the bound state electrons are found. Since the binding energy can be determined with high accuracy, the wave functions pass the test for reproducing the binding energies with a good accuracy. In the next step computer is at work. Sometimes there is a possibility to try the influence of the final state interactions. The usual choices are: plane waves, Coulomb functions...

About twenty years ago such approach was justified to some extent. Most calculations were connected with the characteristics, which could be detected in the experiments. In the real experiments the targets obtained the energies of the order of the electron binding energies. There was no small parameter, and thus no possibility to evaluate the equations, determined by the original

formalism of quantum mechanics. The part of atomic physics, related to the real experiments was indeed becoming the science about the precise computations. The near-threshold behavior of the processes, where one could find a small parameter, was rather an exception.

The situation changed at the end of 80-th, when new synchrotron sources of the photons became available. The experiments with the photon energies of about 10 keV went on. For the light atoms these energies are much larger than the binding energies. A small parameter emerged in the theoretical problems, which could be of practical importance.

The general approach to the "high energy atomic physics" can be based on the works of Bethe [1]. A somewhat modernized presentation is given in [2, 3]. Being initially formulated for the electron-atomic scattering, these principles can be expanded to any process of the interaction between the high energy particles with the bound systems. The wording "high energy" means that the energy exceeds strongly the binding energy of the system. The cross-sections of such processes can be expressed through certain parameters of the bound systems. The interactions between the fast and slow participants can be treated as perturbation. Each act of transferring of large momentum leads to a parametrically small factor. In the region, where the processes are kinematically allowed for the free electrons, a small momentum q of the order of the average momentum of the bound electron η is transferred to the nucleus. The processes outside this kinematical region (which we shall refer to as *Bethe surface*) require large momentum $q \gg \eta$ to be transferred to the nucleus. The processes outside the Bethe surface are thus strongly quenched.

Hence, the amplitudes of the high energy atomic physics processes depend on the parameter

$$\kappa = \eta/q . \quad (1)$$

On the Bethe surface κ is of the order of unity, while $\kappa \ll 1$ outside the Bethe surface. Using the system of units with $\hbar = c = 1$, we can write for the lowest state of the hydrogenlike atom

$$\eta = m\alpha Z , \quad (2)$$

with m being the electron mass, $\alpha = 1/137$ is the fine structure constant, while Z is the charge of the nucleus. We shall use this relation for estimations in the many-electron cases as well.

The perturbative treatment of the fast electrons, which is another point of the Bethe theory, leads to certain misunderstandings sometimes. The sections

"Perturbation theory" of the textbooks on quantum mechanics often treat only the short-ranged forces. Indeed, iteration series term of the Lippman-Schwinger equation for the electron wave function in the Coulomb field of the nucleus contains the infrared-divergent terms. In this sense, there is no such thing as "perturbative series for the Coulomb field" and the words "the fast electrons can be described by the plane waves" become confusing. However, the solution was found long ago. It was guessed by Dalitz [4] and proved by Gorshkov [5], that the divergent terms compose the factor $\exp(i\Phi)$ with real Φ in the amplitudes, thus cancelling in the cross-sections. Hence a way to solve the problem is to present the electromagnetic field of the nucleus in the form $V(r) = -\alpha Z e^{-\lambda r}/r$ with $\lambda \rightarrow 0$. Now the Lippman-Schwinger equation and the perturbative theory become legitimate, while the terms which depend on λ (i.e. $\ln\lambda$) cancel in the cross-sections.

The perturbation parameter of the nonrelativistic electron with the three-dimensional momentum p , moving in the Coulomb field is [6]

$$\xi_Z = m\alpha Z/p. \quad (3)$$

The condition $\xi_Z \ll 1$ means that the relative contribution provided by this interaction to the electron wave function is much smaller than unity [6]. Since different regions of distances may be important for different terms of the expansion of the wave function, one should be careful and expand the *amplitude* in powers of ξ_Z . The small distances are connected with transferring of large momentum $q \gg \eta$ to the nucleus. This process can be considered as perturbation as well, with $\kappa^4 \ll 1$ being the parameter of expansion-Eq(1).

However, one often finds large deviations between the calculations with the plane waves and Coulomb field functions applied for the description of the continuum electrons, even at large values of energies. Sometimes this leads to the statement that "plane wave approximation never works". However, as explained in [7], there are two parameters, which depend on the electron momentum p . Besides the parameter ξ_Z^2 which is the true parameter of the expansion of the wave function at the distances of the order of the size of the atom, there is a parameter $\pi\xi_Z$, which enters through the normalization factor of the Coulomb wave function anyway. In the processes with the large momentum transferred to the nucleus, there are some more terms, containing the parameter $\pi\xi_Z$. Fortunately, they also compose a factor. Thus, assuming

$$\pi\xi_Z \approx 1, \quad \xi_Z^2 \ll 1 \quad (4)$$

one can carry out expansion in powers of ξ_Z^2 . Also, the factors depending on the parameter $\pi\xi_Z$ cancel in the ratios of the cross sections of the processes

containing the high energy electrons with the close values of energies. The recent calculations, presented in [8] illustrated the general validity of the latter statement.

The attempts to include the final state interactions of the outgoing electrons in a straightforward way also met some difficulties, caused by the long-range nature of the interaction. The approach which enabled to include the $e - e$ interaction in the lowest order of the parameter of their interaction

$$\xi^2 = (m\alpha)^2/p_r^2, \quad (5)$$

with p_r being the linear momentum of the relative motion was worked out in [9].

Since the high-energy experiments were not available in atomic physics for a long time, these ideas were not applied much. The only group which systematically used and developed the approach was that of V. G. Gorshkov and his collaborators [10], who focused on the Coulomb field calculations.

Now one can conclude that it is not enough to require that the bound state wave function provides an accurate energy value. It should provide the adequate description of the electron density on the distances, which are important for the considered process. If the process is not allowed for the free electrons, the important distances can be of minor importance for determination of the binding energy. In treating the final states one can use the perturbative expansions in powers of ξ_Z^2 , ξ^2 and κ^2 , taking care that all the terms of the leading order are picked in the amplitude.

It looks to me, that the story of investigation of the double photoionization of helium provides good illustrations of how the application of this approach enabled to obtain the results. It illustrates also, how ignoring of these ideas led to the erroneous results. The theory gave two lessons: at the end of 60-th and at the end of 90-th. The relativistic aspects of the problems will not be touched. I will show, how the main results can be obtained by the standard methods of nonrelativistic quantum mechanics. I will show also, that the Lippman- Schwinger equation appears to be a powerful tool. However, I start with the example of a static problem of the calculation of the ground state binding energy of the helium atom.

2 Calculation of the ground state energy of helium atom

The binding energy of the ground state of helium atom is known from the experiments with very high precision. The relative accuracy was 10^{-6} in late 50-th [2]. It is about $2 \cdot 10^{-7}$ today [11]. This stimulated the attempts to solve the Schrödinger equation for the helium ground state as precisely as possible.

The Hartree-Fock (HF) method, developed in early 30-th, reproduces the value of the binding energy with the relative error of about 1.5%. The experiments are much more precise. The reasons of this failure are known. The HF function does not depend on the distance between the electrons. The probability to find the two electrons to be close to each other is thus overestimated in HF approach.

An alternative approach was developed in the same years as the HF method. It consists in employing approximate wave functions $\psi_a(r_1, r_2, r_{12})$, presented by explicit analytical formulas. Here r_1, r_2 denote the distances between the electron and the nucleus, r_{12} is the distance between the electrons. The lower index a stands for "approximate". These functions contain a number of parameters. The values of the parameters are chosen to minimize the expectation value

$$E = \langle \psi_a | H | \psi_a \rangle , \quad (6)$$

of the Hamiltonian H of the helium atom. The simplest example is the product of two hydrogenlike functions in the field of the nucleus with certain "effective nuclear charge" Z_e . The effective charge plays the role of the variation parameter. In this case

$$\psi_a(r_1, r_2) = \psi_h(r_1)\psi_h(r_2), \quad (7)$$

with the well known analytic expression for $\psi_h(r) = e^{-ar}$, $a = m\alpha Z_e$. (All the wave functions are normalized to $\psi(0) = 1$). The solution $Z_e = 27/16$ [6] provides the energy value with the error of 2%.

More complicated functions of the form

$$\psi_a(r_1, r_2) = e^{-a(r_1+r_2)} P(r_1, r_2, r_{12}) \quad (8)$$

with a and the coefficients of the polynomial

$$P(r_1, r_2, r_{12}) = \sum c_{ijk} r_1^i r_2^j r_{12}^k \quad (9)$$

being the variation parameters, have been used since late 20-th. The earlier versions contained 3 parameters. By the end of 50-th the functions with more than 200 parameters were employed [2]. The functions P , containing also the dependence on the ratios $\frac{r_{12}}{(r_1+r_2)}$ and $\frac{(r_1-r_2)^2}{r_{12}^2}$ have been used. This enabled to obtain the convergence of the results to a certain value, interpreted as the true energy value, with the precision of several units of 10^{-10} , reproducing 5 signs in the value of the binding energy (The calculations with the accuracy better than 10^{-4} require the account of the finite mass and motion of the nucleus, inclusion of relativistic corrections, etc).

However, Barlett *et al.* [12] showed that the functions (8), with P having a polynomial structure could not satisfy the Schrödinger equation for the ground state of helium. Such solution does not exist since it does not have a proper limit at the triple coalescence point $r_1 = r_2 = r_{12} = 0$. The functions which treated the three-particle singularity of the Hamiltonian were found by Fock [13]. They took the form

$$P(r_1, r_2, r_{12}) = \sum c_{ijkn} r_1^i r_2^j r_{12}^k [\ln(r_1^2 + r_2^2)]^n, \quad (10)$$

thus containing the logarithmic terms. (Usually, the Fock function is presented by using the hyperspherical variables [14]).

Inclusion of the logarithmic terms has little effect on the energy value. However, account of these terms improves the convergence of the variation calculations. Also, inclusion of such terms enables to diminish the number of variation parameters. For example, the calculation with the accuracy 10^{-9} required 1078 parameters of the function (9), while it was sufficient to use the function (10) with 52 parameters to achieve the same precision [15]. Furthermore, as noted by Myers *et al.* [16], for some of the functions (9) the variation procedure chooses the parameters in such a way, that the expansion mimicked the function (10) with a smaller number of parameters.

Behavior of the true ground state wave function $\Psi(r_1, r_2, r_{12})$ at the two-particle coalescence points is determined by the Kato cusp conditions [17]. In the point of electron-nucleus coalescence $r_1 = 0$ it is

$$r_0 \frac{\partial \Psi(r_1, r_2, r_{12})}{\partial r_1} = -Z \Psi(0, r_2, r_{12}), \quad (11)$$

while in the electron-electron coalescence point $r_{12} = 0$ it is

$$r_0 \frac{\partial \Psi(r_1, r_1, r_{12})}{\partial r_{12}} = \frac{1}{2} \Psi(r_1, r_1, 0), \quad (12)$$

with $r_0 = \frac{1}{m\alpha}$ standing for the Bohr radius. Equations (11,12) can be viewed as the conditions of the cancellation of the singular terms in the Schrödinger equation.

Let us illustrate this, considering the single-electron wave function $\psi(r)$ of the bound electron in 1s state of certain effective field $U(r)$, which approximates the interactions with the nucleus and with the other electrons. The Schrödinger equation for this case is

$$\frac{-1}{2m} \left(\psi''(r) + \frac{2}{r} \psi'(r) \right) + U(r) \psi(r) = E \psi(r). \quad (13)$$

At $r \rightarrow 0$ the interaction $U(r)$ is determined by interaction with the nucleus. Thus, $U(r \rightarrow 0) \sim -\frac{\alpha Z}{r}$. Hence, for $r \rightarrow 0$ we can write

$$\frac{-1}{2m} \psi''(r) + \lim_{r \rightarrow 0} \left(\frac{\frac{-1}{m} \psi'(r)}{r} - \frac{\alpha Z \psi(r)}{r} \right) = E \psi(r). \quad (14)$$

Since both sides of Eq. (14) should be regular, the expression in brackets should have a finite value. Thus

$$r_0 \frac{\partial \psi(r)}{\partial r} = -Z \psi(0), \quad (15)$$

at $r \rightarrow 0$. Similar analysis of the Schrödinger equation for the atom of helium leads to Eqs.(11),(12). Note, that these relations could have been obtained immediately after the Schrödinger equation was written in 1926. Surprisingly, they were found only thirty years later.

The proper treatment of the two-particle singularity is even more important than the account of the three-particle one [16]. This is because the three particle coalescence point is felt in the smaller volume of integration of the matrix element (6). Myers *et al.* [16] suggested the simplest form of the wave function, which satisfies both Kato conditions

$$\psi_a(r_1, r_2, r_{12}) = e^{-a(r_1+r_2)+br_{12}} \quad (16)$$

with $a = m\alpha Z$, $b = \frac{m\alpha}{2}$. It is amusing that this function, which does not contain fitting parameters, provides the same precision in determination of the binding energy, as the function (7), which contains a variation parameter Z_e . The function (16) knows nothing about the three-particle singularity. The simplest function, which would account both two- and three-particle singularities is described by a more complicated formula, also presented in [16].

If the approximate wave function is chosen in a proper form, the variation procedure exhibits the tendency to find the parameters, providing small errors in the Kato conditions (11,12). The wave functions

$$\psi_a(r_1, r_2, r_{12}) = e^{-a(r_1+r_2)+br_{12}} \sum c_{ijk} r_1^i r_2^j r_{12}^k, \quad (17)$$

($i + j + k \leq N$) analyzed by Teng and Shakeshaft [18] provide the relative errors of about 6% and 25% in Eqs.(11, 12) correspondingly, even in the case of 4 parameters ($N=1$). Although the Kato conditions are satisfied only approximately, it helps in improving the convergence. The function (17) with 14 parameters ($N=4$) provides the value of the binding energy with the relative deviation $3 \cdot 10^{-5}$ from the one, obtained with the functions, described by Eqs (8,9) and containing 210 parameters [2].

In the nowadays studies the subject of investigation is usually not only the expectation value (6), but also the local energy functional

$$E(\vec{r}_1, \vec{r}_2) = \frac{H\psi_a}{\psi_a}, \quad (18)$$

which gives more detailed information about the relation of the approximate function ψ_a to the true wave function Ψ . Another improvement consists in developing the approaches, based on direct solution of the Schrödinger equation. In one of the methods, suggested by Haftel and Mandelzweig (see [19], where the other methods are also discussed) the wave function is presented as a product of a correlation factor, describing the coalescence points singularities and a smooth function, expanded in a series of hyperspherical harmonic functions. The detailed description of these points is beyond the scope of this lecture.

I hope that this Section made one convinced that taking care of the proper analytical structure of the approximate wave functions, one obtains the better results. Now we shall see that this becomes increasingly important for the dynamical problems.

3 Asymptotics of the double photoionization cross section: $\omega^{-7/2}$ or $\omega^{-5/2}$?

The double photoionization of the ground state of helium is studied since the late 50-th. The process provides a direct probe of three-body physics, since the

three charged particles are involved. A huge number of experimental results gives a good possibility for theoreticians to test their approaches.

The mechanism, which determines the main contribution to the nonrelativistic asymptotics of the cross section $\sigma^{2+}(\omega)$ with ω standing for the photon energy, was clarified by Kabir and Salpiter [20]. At the values of ω , exceeding strongly the binding energy I , the process can be viewed as a single photoionization, followed by the transition of the second electron to continuum. The latter process takes place due to the change of the Hamiltonian of the system. This is the analog of the shake-off (*vstryahivanie*), suggested by Feinberg [21] for the description of ionization during the nuclear beta decay, due to the sudden change of the electromagnetic field, caused by the change of the charge of the nucleus. However, using the same wording *shake-off*, one must remember, that, in contrast to the beta decay, the mechanism can not be treated as due to the sudden change of certain effective *field*. Byron and Joachain [22] showed that, being attributed to the sudden change of self-consistent (Hartree-Fock) field, the mechanism underestimates the asymptotic value by a factor of 3. The initial state correlations beyond the effective field are thus of crucial importance.

The electron, which interacts with the photon directly, absorbs most of the energy $\epsilon_1 \approx \omega$. The secondary electron carries the energy of the order of the binding energy $\epsilon_2 \sim I \ll \omega$. Thus, the energy dependence of the amplitude and of the cross section of the high energy nonrelativistic photoionization are the same, as that for the single photoionization process.

Recall now the mechanism of single ionization.

3.1 Asymptotics of the single photoionization

Here we assume that the electrons are described by the single-particle wave functions. The general expression for the amplitude of ionization of $1s$ state is

$$F^+ = (4\pi\alpha)^{1/2} \int d^3r \psi_{p1}^*(r) \gamma e^{i(kr)} \psi(r), \quad (19)$$

with ψ_{p1} being the continuum wave function with the asymptotic momentum $\vec{p}_1$, ψ describes the bound $1s$ electron, while

$$\gamma = -i \frac{(\vec{e} \cdot \vec{\nabla}_r)}{m}, \quad (20)$$

is the operator of interaction between the photon and electron, $\vec{e}$ stands for the photon polarization. The Fourier transform provides

$$F^+ = (4\pi\alpha)^{1/2} \int \frac{d^3q}{(2\pi)^3} \psi_{Fp1}^*(q) \frac{(eq)}{m} \psi_F(q - k), \quad (21)$$

Here the functions ψ_F are the Fourier transforms of the functions ψ . The process requires that momentum

$$\vec{Q} = \vec{k} - \vec{p}_1 \quad (22)$$

is transferred to the recoil ion. Note, that the energy of the fast electron $\epsilon_1 = p_1^2/2m \gg I$, while the binding energy $I = \frac{\eta^2}{2m}$, with η standing for the average momentum of the bound electron. Thus, $\epsilon_1 \gg I$ means, that $p_1 \gg \eta$, i.e. the momentum of the fast electron is much larger than that of the bound electron. Also, the energy dependence of the photon and electron momenta are $k = \omega$ and $p_1 \approx (2m\omega)^{1/2}$. Hence, $Q \gg \eta$ in this process. The large momentum (i.e. momentum, exceeding strongly the average value for the bound electron) should be transferred to the recoil ion.

Start with the plane wave

$$\psi_{Fp1}(\vec{q}) = \psi_{Fp1}^{(0)}(\vec{q}) = (2\pi)^3 \delta(\vec{p}_1 - \vec{q}), \quad (23)$$

for the description of the final state electron. In this case all momentum Q is transferred to the recoil ion by the initial state electron. The contribution to the amplitude is

$$F_0^+ = (4\pi\alpha)^{1/2} \frac{(ep_1)}{m} \psi_F(p_1). \quad (24)$$

For the hydrogenlike functions the asymptotics $\psi_{hF}(p_1) = \frac{8\pi^{-1/2}\eta^{5/2}}{p_1^4} + O(p_1^{-6})$ at $p_1 \gg \eta$ can be obtained explicitly. One can expect the p_1^{-4} behavior to be true in the general case, since the momenta $p_1 \gg \eta$ correspond to small distances $r \ll \eta^{-1}$, where an electron function ψ has a hydrogenlike shape. Anyway, this can be proved rigorously, and the pre-asymptotic coefficient can be calculated. Since the s state wave function does not depend on the direction of the vector $\vec{r}$, we can evaluate

$$\psi_F(p) = \int d^3r \psi(r) e^{-i(\vec{p}\vec{r})} = -\frac{4\pi}{p^2} \int dr r \psi(r) [\cos(pr)]' = \dots - \frac{8\pi\psi'(0)}{p^4} + O(p^{-6}) \quad (25)$$

Here the last equality is obtained after using two integrations by parts.

To get convinced, that Eq.(25) indeed presents the asymptotics of the amplitude, let us try to include the interactions between the outgoing electron and the recoil ion. The lowest order perturbative correction to the outgoing electron wave function, caused by interaction with the nucleus V^C can be obtained by using the standard formula of quantum mechanics

$$\psi_n^{(1)} = \sum \frac{V_{nm}^C \psi_m^{(0)}}{\epsilon_n - \epsilon_m}, \quad (26)$$

with the sum over the states m , which do not coincide with n , $\epsilon_{n,m}$ stand for the unperturbed energy values. This provides

$$\psi_{Fp1}^{(1)}(\vec{q}) = -\frac{8\pi\alpha Zm}{(\vec{p}_1 - \vec{q})^2(p_1^2 - q^2)}, \quad (27)$$

with the contribution

$$F_1^+ = -(4\pi\alpha)^{1/2} \int \frac{d^3q}{(2\pi)^3} \frac{(eq)}{m} \frac{8\pi\alpha Zm}{(\vec{p}_1 - \vec{q})^2(p_1^2 - q^2)} \psi_F(q), \quad (28)$$

to the amplitude. In the amplitude F_1^+ both initial and final state can transfer large momenta q or $p_1 - q$ to the nucleus. The small factor of the order p_1^{-4} is partially compensated by the large phase volume of integration of the order p_1^3 , multiplied by the factor $m\alpha Z$. Thus, the amplitude F_1^+ is indeed ξ_Z times smaller than F_0^+ , dropping faster with the increasing photon energy.

Thus, corrections to asymptotics provide the terms $O(\xi_Z)$, dropping as $\omega^{1/2}$. The large numerical coefficient π can be understood, e.g. by considering the procedure of normalization of the function $\psi_{p1}(r) = \psi_{p1}^{(0)}(r) + \psi_{p1}^{(1)}(r)$. Calculating the value $\psi_{p1}(0) = \psi_{p1}^{(0)}(0) + \psi_{p1}^{(1)}(0) = \int \frac{d^3q}{(2\pi)^3} [\psi_{Fp1}^{(0)}(q) + \psi_{Fp1}^{(1)}(q)]$ one obtains:

$$\psi_{p1}(0) = 1 + \xi_Z J \quad (29)$$

with

$$J = \frac{1}{\pi} \int_0^\infty \frac{dx}{x} \ln \frac{1+x}{|1-x|} = \pi/2, \quad (30)$$

and thus $\psi_{p1}^{(0)} = 1 + \pi\xi_Z/2$. Additional power of π can be viewed as coming from $-i\ln(-1) = \pi$ if the integral J is evaluated in the complex plane. Hence, the amplitude F_0^+ provides the asymptotics of the total amplitude. This leads to the well known asymptotics of the cross section [23]

$$\sigma^+(\omega) = C_1 \omega^{-7/2} \quad (31)$$

Because of the large corrections of the order $\pi\xi_Z$, the nonrelativistic asymptotics can be achieved only for very light atoms. In other cases the relativistic treatment becomes important at the energies far below the nonrelativistic asymptotic region.

3.2 Asymptotics of double photoionization. Spurious controversy

The general expression for the double photoionization amplitude can be written as

$$F^{2+} = \langle \Psi_f | \gamma | \Psi_i \rangle, \quad (32)$$

with Ψ_i and Ψ_f standing for the exact wave functions of the initial and final states, while

$$\gamma = \gamma_1 + \gamma_2; \quad \gamma_j = i \frac{(\vec{e} \vec{\nabla}_j)}{m} \quad (33)$$

describes the interactions of the photon with the two-electron system. The asymptotics of the amplitude [20] is determined by the interactions in the initial state, which transfer small energy $\epsilon_2 \sim I$ to the secondary electron. Large momentum

$$\vec{Q} = \vec{k} - \vec{p}_1 - \vec{p}_2 \quad (34)$$

is transferred to the nucleus.

To obtain the asymptotics, we can describe the final state by the product of the single-particle functions, putting $\Psi_f(\vec{r}_1, \vec{r}_2) = \psi_{p_1}^{(0)}(\vec{r}_1) \psi_{p_2}^C(\vec{r}_2)$. Here ψ^C stands for the continuum wave function in the Coulomb field of the nucleus, while $\psi^{(0)}$ denotes its lowest order of expansion in powers of ξ_Z (the plane wave). Since $p_1 \gg p_2$, we can put $\gamma = \gamma_1$.

Analysis, similar to the one, carried out in the previous subsection, provides

$$F^{2+} = (4\pi\alpha)^{1/2} \frac{(\vec{e} \vec{p}_1)}{m} \int \frac{d^3 q}{(2\pi)^3} \psi_{p_2}^{C*}(\vec{q}) \Psi_F(\vec{p}_1, \vec{q}). \quad (35)$$

The integral on the rhs is saturated by small $q \sim \eta \ll p_1$. The calculation, similar to the one presented by (25), provides $\Psi_F(p_1, q) \sim p_1^{-4}$ at $p_1 \gg \eta$. Thus, at $p_2 \sim \eta$ the dependence of F^{2+} on the photon energy is the same as that of F^+ . The integral over the energy distribution is saturated by $\epsilon_2 \sim I$ and thus does not depend on the photon energy. Hence, the double photoionization

cross section σ^{2+} obtains the same energy dependence as the single ionization cross section, i.e.

$$\sigma^{2+}(\omega) = C_2 \omega^{-7/2} \quad (36)$$

In paper [22] the double photoionization cross section was calculated by employing the product of the Coulomb functions for the description of the final state. The asymptotics was obtained by expansion of the amplitude in powers of ξ_Z . It was noted that the result depends on the form, assumed for the operator of interaction between the photon and electrons. For the *exact* solutions of the Schrödinger equation the velocity form, presented by Eq.(33) is equivalent to the length form

$$\gamma = \gamma_1 + \gamma_2; \quad \gamma_j = i\omega_{fi}(\vec{e}\vec{r}_j) \quad (37)$$

with ω_{fi} being the difference of the energies of the final and initial states, $\omega_{fi} = \omega$ in our case. While the velocity form provided the asymptotics presented by Eq.(36), the length form gave

$$\sigma^{2+}(\omega) = C_2^L \omega^{-5/2}. \quad (38)$$

To understand this, note that since $p_2 \ll p_1$, the important space region is $r_1 \ll r_2$. Thus we can put $\gamma = i\omega(\vec{e}\vec{r}_2)$. Unlike the single photoionization, the electron, which interacts with the photon directly, does not obtain angular momentum. The double photoionization amplitude takes the form

$$F^{2+} = (4\pi\alpha)^{1/2} \omega \int \frac{d^3q_1 d^3q_2}{(2\pi)^6} \psi_{p_1}^{C*}(q_1) \psi_{p_2}^{C*}(q_2) (eq_2) \Psi_F(q_1, q_2). \quad (39)$$

If we describe the fast outgoing electron by the plane wave – Eq.(23), the dependence of the amplitude on the momentum of the fast electron $p_1 \sim (2m\omega)^{1/2}$ is determined by the factor

$$\omega \Psi_F(p_1, q_2) \sim \frac{\omega}{p_1^4}, \quad (40)$$

containing additional energy dependent factor $\omega^{1/2}$, compared to Eq.(24). This leads to Eq.(38) for the asymptotics of the cross section. In the calculations of [22] such behavior remained true, when the fast electron was described by the Coulomb field function.

The authors of [22] suggested that the $\omega^{-5/2}$ result is spurious, and that the corresponding contribution should vanish if the exact wave function are used. This guess was proved by Åberg several years later [24].

3.3 Solution of the controversy

Here I present a simple derivation of the main results of [24]. Large momentum $Q \approx p_1$ can be transferred to the nucleus either by the initial state electron, or by the final state electron. In the length form each of these mechanisms, being treated separately, provides the contribution of the same order in the parametric expansion. The wave function of the outgoing electron can be described by the superposition of the plane wave and the lowest order Coulomb correction. Electron interactions in the final state as well as the higher order Coulomb corrections contribute beyond the asymptotics. The two terms of the expansion of the final state wave function provide the two contributions to the amplitude $F_L^{2+} = F_{L0}^{2+} + F_{L1}^{2+}$, i.e.

$$F_{Lk}^{2+} = (4\pi\alpha)^{1/2} \int \frac{d^3 q_2}{(2\pi)^3} \psi_{Fp2}^*(\vec{q}_2) (\vec{e}\vec{q}_2) A_k(\vec{q}_2) \quad (41)$$

with

$$A_k(\vec{q}_2) = \omega \int \frac{d^3 q_1}{(2\pi)^3} \psi_{Fp1}^{(k)*}(q_1) \Psi_F(\vec{q}_1, \vec{q}_2) \quad (42)$$

and $k = 0, 1$. For the plane wave contribution we find immediately

$$A_0(q_2) = \omega \Psi_F(p_1, q_2), \quad (43)$$

with the large momentum $p_1 \gg \eta$ being transferred to the nucleus by the initial state electrons. (Recall that the lower index F denotes the Fourier transform). The amplitude A_0 is determined by large $q_1 = p_1 \gg \eta$. The integral, corresponding to A_1 is saturated by small momenta $q_1 \sim \eta$. This can be seen immediately from Eq.(27), which presents the function $\psi_{Fp1}^{(1)}$ explicitly. The momentum q_1 can be neglected everywhere, except the wave function Ψ_F . This provides

$$A_1(\vec{q}_2) = -\omega \frac{8\pi\eta}{p_1^4} \int \frac{d^3 q_1}{(2\pi)^3} \Psi_F(\vec{q}_1, \vec{q}_2) = -\frac{8\pi\omega}{p_1^4} \Psi_{PF}(0, q_2). \quad (44)$$

Here we introduced the partial Fourier transform

$$\Psi_{PF}(\vec{r}, \vec{q}_2) = \int d^3 r \Psi(\vec{r}, \vec{r}_2) e^{-i(\vec{q}_2 \vec{r}_2)}. \quad (45)$$

The calculation, similar to that, presented by Eq.(25), gives

$$A_0(q_2) = -\frac{8\pi\omega}{p_1^4} \Psi'_{PF}(0, q_2). \quad (46)$$

Thus, the amplitudes F_{L0}^{2+} and F_{L1}^{2+} appeared to be of the same magnitude. (Recall, that this was not the case for the amplitudes F_0^+ and F_1^+ due to the factor $\frac{(eq)}{m}$ on the rhs of Eq.(28)). This does not contradict to the fact, that the function $\psi^{(1)}(r)$ is ξ_Z times smaller then $\psi^{(0)}(r)$ in any space point. The amplitudes $A_k(q_2)$, which can be presented as

$$A_k(\vec{q}_2) = \omega \int d^3r_1 \psi_{p_1}^{(k)*}(\vec{r}_1) \Psi_{PF}(\vec{r}_1, \vec{q}_2), \quad (47)$$

are dominated by different regions of the space distances r_1 . While A_0 is determined by small $r_1 \sim p_1^{-1}$, much larger distances $r_1 \sim \eta^{-1}$ provide the main contribution to A_1 .

Treated separately, each of the terms $A_{0,1}$ would provide $\omega^{-5/2}$ law for the cross section. However, carrying out the inverse Fourier transform, one finds $A_0 + A_1 = 0$ for any q_2 due to the Kato condition (11), and thus

$$F_{L0}^{2+} + F_{L1}^{2+} = 0. \quad (48)$$

This shows, how the straightforward calculations in the length form, employing the approximate wave functions, which do not satisfy Eq.(11), provide the spurious $\omega^{-5/2}$ asymptotics for the double photoionization cross section.

3.4 The Lippman-Schwinger equation

In this subsection I show that the Lippman-Schwinger equation (LSE) is a convenient tool for the asymptotical analysis. The general form of LSE for the two-electron system is

$$\Psi = \Psi^{(0)} + G^\epsilon V \Psi. \quad (49)$$

Here V stands for the interaction, ϵ denotes the energy value, corresponding to the solution Ψ of the Schrödinger equation, $\Psi^{(0)}$ and G^ϵ are the solution and the Green function for $V = 0$ (free motion).

To simplify the notations, let us write the LSE for the single-particle wave function ψ , which we assume to be the exact solution of the Schrödinger equation in a certain effective field $U(r)$. It is

$$\psi = \psi^{(0)} + G^\epsilon U \psi. \quad (50)$$

Now G^ϵ is the free single-particle Green function. In the momentum space

$$\langle \vec{p} | G^\epsilon | \vec{f} \rangle = \frac{\delta(\vec{p} - \vec{f})}{\epsilon - p^2/2m + i\nu} \quad \nu \rightarrow 0. \quad (51)$$

For the bound states $\psi^{(0)} = 0$ and

$$\psi_F(p) = \frac{2m}{2m\epsilon - p^2} \int \frac{d^3q}{(2\pi)^3} \langle \vec{p} | U | \vec{q} \rangle \psi_F(\vec{q}), \quad (52)$$

with ϵ standing for the single-particle binding energy. In the asymptotics $2m\epsilon \ll p^2$. The integral is saturated by $q \sim \eta$. Thus, Eq.(52) presents the bound state wave function $\psi_F(p)$ at large momenta $p \gg \eta$ in terms of the same wave function at small momenta $q \sim \eta$.

For s states we can put $\langle \vec{p} | U | \vec{q} \rangle = \langle \vec{p} | U | 0 \rangle$. Since large p correspond to small r , only the interactions with the nucleus are important in the asymptotics. Thus, putting $\langle \vec{p} | U | \vec{q} \rangle = -\frac{4\pi\alpha Z}{p^2}$, we obtain

$$\psi_F(p) = \frac{8\pi\alpha Z\psi(0)}{p^4} + O(p^{-6}), \quad (53)$$

Due to the Kato condition (15), this expression is identical to Eq (25). In similar way we can find the asymptotics of the wave function for the ground state of helium

$$\Psi_F(p, q_2) = \frac{8\pi\alpha Z\Psi_{PF}(0, q_2)}{p^4} + O(p^{-6}). \quad (54)$$

with the partial Fourier transform Ψ_{PF} defined by Eq.(45).

As to the wave function of the outgoing electron, the first iteration of the LSE provides the same result, as Eq.(26) of quantum mechanics (adding an imaginary infrared divergent term, which will be absorbed into the phase factor, mentioned in the Introduction). Thus, calculating the double photoionization amplitude in the length form, we immediately find the cancellation, expressed by Eq.(48). Hence, in the LSE technique one would not obtain the spurious $\omega^{-5/2}$ term in the asymptotics of the double photoionization cross section, even being not aware of the Kato conditions.

4 Shape of the double photoionization spectrum: U or W ?

4.1 Mechanisms of the process

The progress of the experimental facilities in early 90-th made possible the measurements of the energy distributions of the double photoionization of he-

lithium. The low energy measurements have been carried out. The results for the high energy are expected soon. This stimulated the theoretical investigations.

Evaluation of the shape of the spectrum curve have been obtained in [25]. The energy distribution $d\sigma^{2+}/d\varepsilon$ reaches its largest values at the edge of the spectrum, where $\varepsilon_2 \sim I \ll \omega$ or $\varepsilon_1 \sim I \ll \omega$. The shake-off mechanism is dominative here [20]. Large momentum $Q \gg \eta$ is exchanged between the nucleus and the fast electron. Considering the larger values $\varepsilon_{1,2} \gg I$, we need both electrons to carry large momenta $p_i \gg \eta$. If the energies of the outgoing electrons are not too close, the process can be viewed as the single photoionization, followed by the electron scattering [26, 27]. This *final state scattering* (FSS) mechanism dominates until we do not reach the very vicinity of the center of the spectrum, where the *quasifree* (QF) process becomes possible.

To understand the point, recall that each act of the exchange by a large momentum leads to a small factor in the amplitude. The FSS mechanism requires the exchange by two large momenta. However in the vicinity of the center we can manage by a single large momentum exchange. The electron, which interacts with the photon directly, can exchange large momentum with the second electron without participation of the nucleus. Unlike the single photoionization, the process can go on with the free electrons. The Bethe surface is determined by the condition $\vec{Q} = 0$, *i.e.*

$$\vec{p}_1 + \vec{p}_2 = \vec{k}, \quad (55)$$

which requires

$$\delta = \frac{\varepsilon_1 - \varepsilon_2}{\omega} \leq \sqrt{\frac{\omega}{\omega + m}} \ll 1. \quad (56)$$

(Recall, that we do not consider very large values of the photon energies $\omega \sim m$, which require the relativistic treatment of the outgoing electrons.) In other words, the free kinematics becomes possible in the vicinity of the center of the spectrum.

Following the general ideas, presented in the Introduction, we expect the surplus of the spectrum curve near it's central point due to the described QF mechanism. The energy distribution is dominated by small values $Q \sim \eta$ of the momentum, transferred to the nucleus.

To calculate the asymptotics, we can neglect the terms of the order Q/p_i in the wave function of the outgoing electrons. The QF amplitude can be presented as

$$F_{QF}(\vec{p}_1, \vec{p}_2) = F_{QF}(\vec{p}, \vec{Q}) = D(Q^2)F_{fr}(\omega, \delta) \quad (57)$$

with $\vec{p} = (\vec{p}_1 - \vec{p}_2)/2$. The factor $D(Q^2)$ contains all the properties of the initial state. Its explicit form will be given below — Eq. (71). The amplitude F_{fr} of the absorption of the photon by two free electrons at rest can be presented as

$$F_{fr}(\omega, \delta) = (\vec{e}\vec{p}_1)f(\omega, \delta) + (\vec{e}\vec{p}_2)f(\omega, -\delta) , \quad (58)$$

since the electron system is a space-symmetric state. We do not need to clarify the form of the function $f(\omega, \delta)$ here. Anyway, since $\delta \ll 1$, the lowest order of expansion in powers of δ gives

$$F_{fr}(\omega, 0) = (\vec{e}, \vec{p}_1 + \vec{p}_2)f(\omega, 0) = (\vec{e}\vec{k})f(\omega, 0) = 0, \quad (59)$$

with the lowest order nonvanishing term being

$$F_{fr}(\omega, \delta) = 2(\vec{e}\vec{p})\delta f'(\omega, 0) . \quad (60)$$

Here f' denotes the derivative with respect to δ .

The energy distribution in the vicinity of the center is the interplay of FSS and QF mechanisms. Their contributions change with the photon energy in different ways. At $\omega \ll 2keV$ the FSS dominates, providing **U** shaped curve. At $\omega \approx 2keV$ the QF mechanism becomes more important. The curve obtains **W** shape.

Note that usually the photoionization calculations are carried out in the lowest order of expansion in powers of kr_c with r_c being the characteristic value of the distances, which are important in the process. This is known as the dipole approximation. In the nonrelativistic photoionization $r_c \sim p^{-1} \sim (m\omega)^{-1/2}$ and the dipole approximation corresponds to the lowest order of expansion in powers of $k/p \sim (\omega/m)^{1/2}$. Going beyond the dipole approximation have been something unusual, at least until the recent time. The dipole approximation QF amplitude turns to zero due to Eq. (59). This happens because the two-electron system cannot carry the orbital momentum, equal to unity, in this kinematics. One should go beyond the dipole approximation to obtain the nonvanishing QF amplitude. Unfortunately, this point was not emphasized strongly enough neither in [25], nor in a more detailed paper [27]. The QF amplitude is expressed by Eqs. (57) and (60) with

$$\delta = \frac{(\vec{p}\vec{k})}{m\omega} . \quad (61)$$

4.2 Dipole approximation calculations

Meanwhile, several calculations of the high energy photoionization spectrum have been reported. They were based on the straightforward computation of the amplitude in the dipole approximation. The results appeared to be controversial. In [28] the spectrum curve had **U** shaped form if the final state interactions (FSI) between the outgoing electrons were neglected. Account of the FSI provided **W** shaped curve. The authors of [29] faced the opposite situation. Their spectrum curve for $\omega = 2.8keV$ had **W** shape with a very flat peak, when FSI have been neglected. The computations with FSI included provided **U**-shaped spectrum. Another series of calculations without FSI [30, 31] gave a peak in the center of the energy distribution.

4.3 Explanation of the dipole approximation results

To understand what happened [32, 33], present the ground state wave function in variables $\vec{R} = (\vec{r}_1 + \vec{r}_2)/2$ and $\vec{\rho} = \vec{r}_1 - \vec{r}_2$, introducing $\tilde{\Psi}(\vec{R}, \vec{\rho}) = \Psi(\vec{r}_1, \vec{r}_2)$. Describing the outgoing electrons by the plane waves, we find for the amplitude

$$F^{(0)} = (4\pi\alpha)^{1/2} \int d^3R d^3\rho e^{i(\vec{Q}\vec{R}) - i(\vec{p} - \vec{k}, \vec{\rho})} \gamma \tilde{\Psi}(\vec{R}, \vec{\rho}). \quad (62)$$

Following the analysis, carried out in Sec.3, we expect the asymptotics $F^{(0)}(p, Q) \sim 1/(p^4 Q^4)$ at $p, Q \gg \eta$. However, in the vicinity of the center of the energy distribution we can make $Q \sim \eta$. These values of Q will be dominative in the energy distribution. The calculation, similar to the one, presented by Eq. (25) provides

$$F^{(0)} = -(4\pi\alpha)^{1/2} \frac{(\vec{e}\vec{Q})}{m} \cdot \frac{8\pi\alpha}{p^4} \int d^3R e^{i(QR)} r_0 \frac{\partial \tilde{\Psi}(R, \rho=0)}{\partial \rho} \quad (63)$$

at $p \gg \eta$, $Q \sim \eta$, with $\partial \tilde{\Psi}(R, \rho=0)/\partial \rho$ denoting the partial derivation of the function $\tilde{\Psi}(R, \rho)$ with respect to ρ at $\rho = 0$. The energy dependence of the amplitude $F^{(0)}$ is determined by the factor p^{-4} , caused by the large momentum exchange.

Large momentum can be exchanged by the outgoing electrons as well. The lowest order perturbative correction $\psi^{(1)}$ to the final state wave function can be obtained by using the standard formula (26) with V^C being replaced by

the interaction between the electrons. The corresponding amplitude $F^{(1)}$ is of the same order of magnitude as $F^{(0)}$, containing the same factor p^{-4} . The situation is similar to that with the shake-off amplitude in the length form, which was discussed above. The correction $\psi^{(1)}(\rho)$ to the wave function of the outgoing electrons is ξ times smaller than the free wave function $\psi^{(0)}(\rho)$ in each space point ρ . However, the functions $\psi^{(0)}$ and $\psi^{(1)}$ contribute to the amplitude in different space regions. While the amplitude $F^{(0)}$ is determined by small distances $\rho \sim p^{-1}$, the amplitude $F^{(1)}$ is dominated by the values of ρ of the order of the size of the atom $\sim \eta^{-1}$. In both cases $R \sim Q^{-1} \sim \eta^{-1}$, and both electrons are separated from the nucleus by the distances of the order of the size of the atom.

Direct calculation provides

$$F^{(1)} = (4\pi\alpha)^{1/2} \frac{(\vec{e}\vec{Q})}{m} \cdot \frac{4\pi\alpha}{p^4} \int d^3R e^{i(QR)} \tilde{\Psi}(R, \rho = 0) \quad (64)$$

at $p \gg \eta$, $Q \sim \eta$. Each of the amplitudes $F^{(0,1)}$, treated separately, provide the contribution, which exceeds strongly the amplitude, describing the FSI mechanism and requiring $Q \gg \eta$. Hence, each of the terms $F^{(0,1)}$, treated separately, would lead to a surplus in the energy distribution in the central region. However

$$\begin{aligned} F^{(0)} + F^{(1)} &= (4\pi\alpha)^{1/2} \frac{(\vec{e}\vec{Q})}{m} \frac{4\pi\alpha}{p^4} \int d^3R e^{i(QR)} \\ &\times \left(\tilde{\Psi}(R, \rho = 0) - 2r_0 \frac{\partial \tilde{\Psi}(R, \rho = 0)}{\partial \rho} \right) = 0, \end{aligned} \quad (65)$$

since the last factor in the integrand turns to zero for all values of R due to Eq. (12). In the dipole approximation the leading contribution to the energy distribution is determined by $Q \gg \eta$, providing the **U**-shaped spectrum curve. However, the true distribution can be obtained only by going beyond the dipole approximation.

Now we can explain the results of [28]–[31]. The authors of [28] used the ground state wave function with a weak dependence on ρ . It was too weak to provide a maximum, when neglecting FSI. Inclusion of FSI provided a spurious uncompensated **W** peak. In contrast, authors of [29] used wave functions which satisfied the Kato condition (12). Thus they obtained **U** shape, when taking into account the FSI. This corresponds to Eq. (65). However, a spurious **W** peak emerged when the FSI were neglected. The spurious central surplus,

obtained in [30, 31] in dipole approximation with FSI being neglected have the same origin.

The Lippman-Schwinger equation (LSE), presented by Eq. (49) enables to obtain these results in a simple way. The amplitude $F^{(0)}$ can be expressed in terms of asymptotics of the Fourier transformed wave function

$$F^{(0)} = -(4\pi\alpha)^{1/2} \frac{(\vec{e}\vec{Q})}{m} \tilde{\Psi}_F(-\vec{Q}, \vec{p}) , \quad (66)$$

while the asymptotics of the function $\tilde{\Psi}_F(\vec{p}, \vec{Q})$ is provided by LSE equation

$$\tilde{\Psi}_F(-\vec{Q}, \vec{p}) = -\frac{4\pi m\alpha}{p^4} \int \frac{d^3q}{(2\pi)^3} \tilde{\Psi}_F(-\vec{Q}, \vec{q}) \quad (67)$$

at $Q \sim \eta$, $p \gg \eta$. The integral in the rhs is saturated by $q \sim \eta$. Hence, the LSE presented the function $\tilde{\Psi}_F(Q \sim \eta, p \gg \eta)$ in terms of this function at $Q \sim \eta$, $q \sim \eta$. Using Eqs. (64) and (67) we immediately obtain the cancellation $F^{(0)} + F^{(1)} = 0$, without employing the Kato condition (12).

Note that the interactions between the outgoing electrons appeared to be more important than their interactions with the nucleus. The latter provide only additional corrections, while the former determine a possible mechanism of the process. Similar analysis with the Coulomb functions for the continuum electrons as a starting point is given in [32].

4.4 Necessary properties of the approximate functions

Most of the calculations of the energy spectrum are based on the original formalism of quantum mechanics. This means straightforward employing of the equation

$$d\sigma^{2+} = \frac{\pi}{\omega} |\bar{F}|^2 \delta(\varepsilon_1 + \varepsilon_2 - \omega - I^{2+}) \frac{d^3p_1}{(2\pi)^3} \frac{d^3p_2}{(2\pi)^3} \quad (68)$$

with the amplitude F calculated as

$$F = \int d^3r_1 d^3r_2 \psi_{fa}^*(\vec{r}_1, \vec{r}_2) \left(\gamma_1 e^{i(\vec{k}\vec{r}_1)} + \gamma_2 e^{i(\vec{k}\vec{r}_2)} \right) \psi_a(\vec{r}_1, \vec{r}_2) \quad (69)$$

by the direct calculation of the integral. One should inevitably use *approximate* wave functions ψ_{fa} and ψ_a for the description of the final and initial states.

To reproduce essential physics, the approximate wave functions should possess certain properties of the true wave functions.

As we have seen in Sec. 3, in the region of the shake-off domination $\varepsilon_1 \approx \omega$, $\varepsilon_2 \sim I$, the FSI correlations are not important and one can approximate the final state by the product of the single-particle functions, putting $\psi_{fa}(\vec{r}_1, \vec{r}_2) = \psi_{ap_1}(r_1)\psi_{ap_2}(r_2)$ and employing the Coulomb field function for ψ_{ap_2} . In the velocity form of the operator γ one can use the plane wave for the description of the fast electron [20]. Description of the initial state by the product of the single-particle wave functions

$$\psi_a(\vec{r}_1, \vec{r}_2) = \psi_s(r_1)\psi_s(r_2) \quad (70)$$

leads to the quantitatively wrong results. Thus the initial state correlations beyond effective field approximation are important [22]. However, there is no formula, which would enable to select "good" wave functions. One should be even more careful, employing the length form of the operator γ . The lowest order Coulomb correction to the wave function of the fast outgoing electron should be included and Eq. (11) should be fulfilled for the initial state function $\psi_a(\vec{r}_1, \vec{r}_2)$ [24].

Turning to the part of the spectrum with $\varepsilon_{1,2} \gg I$ we see that the lowest order final state interactions should be included in the wave function ψ_{fa} . This is sufficient for the proper description of all the region of FSS domination. However, to describe the region of QF mechanism domination, one needs the wave function ψ_a , for which the Kato condition (12) is true. Otherwise the spectrum curve obtains a spurious **W** peak in the dipole approximation (the latter corresponds to replacement $e^{i(kr_{1,2})}$ by unity in Eq. (69)).

As we saw earlier, the true **W** peak can be obtained beyond the dipole approximation. This is equivalent to evaluation of the rhs of Eq. (69), by including the second terms of the expansion of the exponential factors $e^{i(\vec{k}\vec{r}_j)} = 1 + i(\vec{k}\vec{r}_j) + \dots$. Such approach provides Eq. (57) for the QF amplitude with $F_{fr}(\omega, \delta)$ defined by Eq. (60) and

$$D(Q^2) = 2r_0 \int d^3R e^{i(QR)} \frac{\partial \tilde{\psi}_a(R, \rho=0)}{\partial \rho}. \quad (71)$$

The proper value of the derivative $\frac{\partial \tilde{\psi}_a(R, \rho=0)}{\partial \rho}$ can be insured by the Kato condition (12). Thus, even taking care of separating the quadrupole terms, one needs the approximate function $\tilde{\psi}_a$, satisfying the Kato condition (12) for the

proper description of the QF central peak. Of course, any approximate function with a nonzero value of $\frac{\partial \tilde{\psi}_a(R, \rho=0)}{\partial \rho}$ provides a central peak describing the QF mechanism qualitatively. However, the quantitative results can be trusted only if Eq.(12) is satisfied. Considering such approximate function, which can be presented as combinations of the single-particle functions (70), we find the corresponding

$$\tilde{\psi}_a(\vec{R}, \vec{\rho}) = \tilde{\psi}_s\left(\vec{R} + \frac{\vec{\rho}}{2}\right) \tilde{\psi}_s\left(\vec{R} - \frac{\vec{\rho}}{2}\right) \quad (72)$$

to be the even function of ρ . Thus, the computations, carried out in the quadrupole approximation would show no trace of the QF contribution. The Hartree-Fock functions provide one of examples.

5 Summary

We saw that theoretical analysis, preceding the computer calculations, appears to be very helpful. In the static problem of the calculation of the ground state energy of helium atom it improves the convergence of the procedure and enables to diminish the number of phenomenological parameters. It helps to avoid "submarine rocks" in the calculations of the asymptotics of the double photoionization cross section. It enables also to find the true shape of the double photoionization spectrum curve.

It was shown also that the Lippman-Schwinger equation is a powerful tool for investigation of the asymptotics. Unfortunately, the LSE and the *Feynman diagram technique*, which can be viewed as its visual illustration, did not yet become a part of everyday life in atomic theory studies.

I am indebted to M.Ya. Amusia, A.I. Mikhailov, R.H. Pratt, T. Surić and especially to V.G. Gorshkov for helpful discussions.

References

- [1] H. A. Bethe, in *Handbuch der Physik*, Springer, Berlin, 1933, v.24/1.
- [2] H. A. Bethe and R. W. Jackiw, *Intermediate Quantum Mechanics*, Benjamin, NY, 1968.
- [3] M. Inokuti, Rev. Mod. Phys. **43**, 297 (1971).

- [4] R. H. Dalitz, Proc. R. Soc. A **206**, 509 (1951).
- [5] V. G. Gorshkov, JETP, **40**, 1481 (1960).
- [6] L. D. Landau and E. M. Lifshitz, *Quantovaya mekhanika*, Nauka, Moscow 1974-in Russian; *Nonrelativistic Quantum Mechanics*, Pergamon, Oxford, 1974.
- [7] V. G. Gorshkov, A. I. Mikhailov and V. S. Polikanov, Nucl. Phys. **55**, 273 (1964).
- [8] N. B. Avdonina, E. G. Drukarev and R. H. Pratt, Phys. Rev. A **65**, 052705 (2002).
- [9] E. G. Drukarev and M.I. Strikman, Phys. Lett. B **186**, 1 (1987).
- [10] V. G. Gorshkov, *Materialy VII Zimnei Shkoly LNPI-* in Russian, *Proceedings of VII LNPI Winter School*, **2**, 415 1972.
- [11] K. S. E. Eikema, W. Ubachs, W. Vassen, and W. Hogervost, Phys. Rev. Lett, **71**, 1690 (1993).
- [12] J. H. Barlett, J. J. Gibbons and G. G. Dunn, Phys. Rev. **47**, 679 (1935).
- [13] V. A. Fock, Izv. Akad. Nauk SSSR, Ser Fiz. **18**, 161 (1954)-in Russian; K.Nor. Vidensk. Selsk. Forh. **31**, 138, 145 (1958).
- [14] M. G. Veselov and L. N. Labzovskii, *Teoriya Atoma. Stroenie Electronnyh Obolochek*, Nauka, Moscow 1986-in Russian.
- [15] A. M. Ermolaev, Vest. LGU, ser. Fyz. **14**, 46 (1958); **16**, 19 (1961).
A. M. Ermolaev and G. B. Sochilin, Dokl. Akad. Nauk SSSR **155**, 1050 (1964)-in Russian; Soviet Phys.-Doklady **9**, 292 (1964).
G. B. Sochilin, Int. J. Quant. Chem., **3**, 297 (1969).
- [16] C. R. Myers, C. J. Umrigar, J. P. Sethna, and J. D. Morgan III, Phys. Rev A **44**, 5537 (1991).
- [17] T. Kato, Comm.Pure. Appl. Math. **10**, 151 (1957).
- [18] Z. Teng and R. Shakeshaft, Phys. Rev. A **47**, R3487 (1993).
- [19] M. I. Haftel and V.B. Mandelzweig, Ann. Phys. **189**, 29 (1989).

- [20] P. K. Kabir and E. E. Salpiter, Phys. Rev. **108**, 1256 (1957).
- [21] E. L. Feinberg, Dokl. Akad. Nauk SSSR **23**, 778 (1939)- in Russian; Journ. of Phys. **4**, 423 (1941).
A.B. Migdal, Journ. of Phys. **4**, 449 (1941).
- [22] F. W. Byron Jr. and C. J. Joachain, Phys. Rev. **164**, 1 (1967).
- [23] H. Bethe, and E. E. Salpiter, *Quantum Mechanics of One-and Two-Electron Atoms*, Springer-Verlag, Berlin, 1958.
- [24] T. Åberg, Phys.Rev. A **2**, 1726 (1970).
- [25] M. Ya. Amusia, E. G. Drukarev, V. G. Gorshkov, and M. P. Kazachkov, J. Phys. B **8**, 1248 (1975).
- [26] E. G. Drukarev, V. G. Gorshkov, A. I. Mikhailov, and S. G. Sherman, Phys.Lett. A **46**, 467 (1974).
- [27] E. G. Drukarev, Phys. Rev. A **52**, 3910 (1995).
- [28] M. A. Kornberg and J. E. Miraglia, Phys. Rev. A **48**, 3714 (1993).
- [29] Z. Teng and R. Shakeshaft, Phys. Rev. A **49**, 3597 (1994).
- [30] M. A. Kornberg and J. E. Miraglia, Phys. Rev. A **60**, R1743 (1999).
- [31] M. A. Kornberg and J. E. Miraglia, Eur. Phys. J. D **12**, 45 (2000).
- [32] T. Surić, E. G. Drukarev and R. H. Pratt, Phys. Rev. A **67**, 022709 (2003).
- [33] E. G. Drukarev, N. B. Avdonina and R. H. Pratt, J. Phys. B **34**, 1 (2001).